



Rhode Island Pediatric State-Supplied Vaccines¹

Vaccine Type	Brand Name	Rhode Island Policy Guidelines For Vaccine Use ^{2,3}	CPT Code	Dose	Route	Needle Size	Manufacturer	Thimerosal Concentration ⁴
RECOMMENDED VACCINES								
DTaP-Hep B-IPV combination	Pediarix	Routine: primary series at 2, 4, & 6 months	90723	0.5 mL	IM	22-25g 1-1½" ⁿ	GlaxoSmithKline 1-866-475-8222	Free
DTaP (Diphtheria-tetanus-pertussis)	Infanrix	Routine: 4th dose at 15-18 months	90700	0.5 mL	IM	22-25g 1-1½" ⁿ	GlaxoSmithKline 1-866-475-8222	Free
DTaP-IPV (Polio) combination	Kinrix	Routine: 5th dose DTaP & 4th dose IPV at 4-6 years	90696	0.5 mL	IM	22-25g 1-1½" ⁿ	GlaxoSmithKline 1-866-475-8222	Free
Hepatitis A	Havrix	Routine: 2 doses 6 months apart to children 12-23 months & high-risk 2-18 yrs Permissive: 2-18 years of age	90633	0.5 mL	IM	22-25g 1-1½" ⁿ	GlaxoSmithKline 1-866-475-8222	Free
Hepatitis B	Engerix B	Routine: birth dose; catch-up vaccination through 18 years of age	90744	0.5 mL	IM	22-25g 1-1½" ⁿ	Merck 1-800-637-2579	Free
Hib (Haemophilus influenzae type B)	PedvaxHIB	Routine: 3 doses at 2, 4, & 12-15 months	90647	0.5 mL	IM	22-25g 1-1½" ⁿ	Sanofi Pasteur 1-800-822-2463	Free
HPV4 Human Papillomavirus	Gardasil	Females- Routine: 11-12 years; 3 doses at 0, 1-2, 6 month schedule; catch-up vaccination through 18 years Males- Permissive: 9-18 years of age; 3 doses at 0, 1-2, 6 month schedule	90649	0.5 mL	IM	22-25g 1-1½" ⁿ	Merck 1-800-637-2579	Free
Influenza ⁵	Fluzone (Pediatric)	Routine: children 6-35 months	90655	0.25 mL	IM	22-25g 1-1½" ⁿ	Sanofi Pasteur 1-800-822-2463	Free
	Fluzone	Routine: children 3-18 years	90656	0.5 mL				
	Afluria	Routine: children 3-18 years	90658	0.5 mL			Merck 1-800-637-2579	Free
	FluMist	Routine: healthy children 2-18 years	90660	0.2 mL			Nasal Spray	N/A
MCV4 (Meningococcal conjugate)	Menactra	Routine: 1 dose at 11-12 yrs; catch-up 13-18 yrs; high-risk children 2-10 years	90734	0.5 mL	IM	22-25g 1-1½" ⁿ	Sanofi-Pasteur 1-800-822-2463	Free
MMR (Measles-mumps-rubella)	MMRII	Routine: 2 doses at 12-15 months and 4-6 years	90707	0.5 mL	SC	23-25g 5/8" ⁿ	Merck 1-800-637-2579	Free
PCV13 (Pneumococcal conjugate)	Prenvar	Routine: 4 doses at 2, 4, 6, 12-15 months; 5 th dose only for children < 5 years if all previous doses were PCV7	90670	0.5 mL	IM	22-25g 1-1½" ⁿ	Wyeth Lederle 1-800-572-8221	Free
RV (Rotavirus)	Rotarix	Routine: 2 doses at 2 & 4 months	90681	2.0 mL	PO (Oral)	NA	GlaxoSmithKline 1-866-475-8222	Free
Tdap (Tetanus-diphtheria-pertussis)	Boostrix	Routine: 1 dose at 11-12 years; catch-up vaccination through 18 years	90715	0.5 mL	IM	22-25g 1-1½" ⁿ	GlaxoSmithKline 1-866-475-8222	Free
Varicella (Chickenpox)	Varivax	Routine: 2 doses at 12-15 mos & 4-6 yrs; Catch-up 2 nd dose through 18 years	90716	0.5 mL	SC	23-25g 5/8" ⁿ	Merck 1-800-637-2579	Free
VACCINES USED IN SPECIAL CIRCUMSTANCES								
DT (Diphtheria-tetanus)	DT	Use for children < 7 years with contraindication to pertussis vaccine	90702	0.5 mL	IM	22-25g 1-1½" ⁿ	Sanofi Pasteur 1-800-822-2463	Trace < 0.00012%
Td (Tetanus-diphtheria)	Decavac	Use for children 7-9 years and > 10 years for catch-up	90714	0.5 mL	IM	22-25g 1-1½" ⁿ	Sanofi Pasteur 1-800-822-2463	Trace < 0.00012%
IPV (Polio)	IPOL	Use for children > 6 years for catch-up	90713	0.5 mL	IM	22-25g 1-1½" ⁿ	Sanofi Pasteur 1-800-822-2463	Free
PPV23 (Pneumococcal polysaccharide)	Pneumovax	Use for children 2-18 years at high risk for pneumococcal disease	90732	0.5 mL	SC	23-25 g 5/8" ⁿ	Merck 800-637-2579	Free

¹ Pediatric vaccine is provided to Rhode Island healthcare providers delivering care in RI for children (up to 19 years of age) through Rhode Island's Universal Vaccine Program.

² Routine vaccine recommendations: Vaccine should be proactively provided as the standard of care to all children up to 19 years of age.

³ Permissive vaccine recommendations: Vaccine should be provided upon parental request; the provider may proactively offer the vaccine during office visits.

⁴ Source: US Food and Drug Administration, Table 3: Thimerosal and Expanded List of Vaccines: www.fda.gov/cber/vaccine/thimerosal.htm

⁵ Children < nine years of age receiving influenza vaccine for the first time need two doses separated by at least four weeks.